

Fundamental solutions of homogeneous elliptic differential operators.

Brice Camus

*Ruhr-Universität Bochum, Fakultät für Mathematik,
Universitätsstr. 150, D-44780 Bochum, Germany.
Email : brice.camus@univ-reims.fr*

Abstract

We compute fundamental solutions of homogeneous elliptic differential operators, with constant coefficients, on $\mathbb{R}^n$ by mean of analytic continuation of distributions. The result obtained is valid in any dimension, for any degree and can be extended to pseudodifferential operators of the same type.

Key words: Fundamental solutions; PDE.

Let be $P := P(D_x)$ a pseudodifferential operator, with constant coefficients, obtained by mathematical quantization of the function p , i.e :

$$Pf(x) = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{i\langle \xi, x \rangle} p(\xi) \hat{f}(\xi) d\xi. \quad (1)$$

Here, and in the following, the notation :

$$\hat{f}(\xi) = \int_{\mathbb{R}^n} e^{-i\langle y, \xi \rangle} f(y) dy,$$

designs the Fourier transform of f . We note $\mathcal{S}(\mathbb{R}^n)$ the Schwartz space and $\mathcal{S}'(\mathbb{R}^n)$ the distributions on $\mathcal{S}(\mathbb{R}^n)$. A fundamental solution for P is a distribution $\mathfrak{S} \in \mathcal{S}'(\mathbb{R}^n)$ such that $P\mathfrak{S} = \delta$. Fundamental solutions play a major role in the theory of PDE. For a large overview on this subject, and applications, we refer to [1,2]. Apart in some trivial cases, there is few explicit characterizations of fundamental solutions and most results concern the existence. The case of order 3 homogeneous operators, in dimension 3, was treated in [3].

¹ Work supported by the *SFB/TR12* project, *Symmetries and Universality in Mesoscopic Systems*.

Always for $n = 3$, the case of certain elliptic homogeneous operators of degree 4 was also solved in [4]. We are here interested in the case of a definite homogenous polynomial p_k on $\mathbb{R}^n$, i.e. $p_k(\xi) = 0 \Leftrightarrow \xi = 0$ and :

$$p_k(\lambda\xi) = |\lambda|^k p_k(\xi).$$

Note that k has to be even but we do not impose the spherical symmetry of p_k . Strictly speaking, it is not necessary to assume that $k \in \mathbb{N}$ and we can consider operators with a conical singularity at the origin. The main motivation is that such an operator can generalize the Laplacian (positivity and ellipticity) but these operators also play a role in physical optics. To state the main result, we introduce the spherical average of $g \in \mathcal{S}(\mathbb{R}^n)$ w.r.t. the symbol p :

$$A(g)(r) = r^{n-1} \int_{\mathbb{S}^{n-1}} g(r\theta) p_k(\theta)^{-1} d\theta. \quad (2)$$

Here $p_k(\theta)$ designs the restriction of p_k on the sphere, which defines a strictly positive function. Clearly, we have $A(g) \in \mathcal{S}(\mathbb{R}_+)$ and we can extend $A(g)$ by 0 for $r < 0$ to obtain a L^2 function on the line. The main result is :

Theorem 1 *If $k < n$ a fundamental solution $\mathfrak{S} \in \mathcal{S}'(\mathbb{R}^n)$ for P is given by :*

$$\langle \mathfrak{S}, f \rangle = \frac{1}{(2\pi)^n} \int_{\mathbb{R}_+ \times \mathbb{S}^{n-1}} p_k(\theta)^{-1} \hat{f}(r\theta) r^{n-1-k} dr d\theta.$$

But, when $k \geq n$, we have :

$$\langle \mathfrak{S}, f \rangle = -C_{k,n} \frac{\partial^{k-1} A(\hat{f})}{\partial r^{k-1}}(0) + D_{k,n} \int_{\mathbb{R}_+} \log(u) \frac{\partial^k A(\hat{f})}{\partial r^k}(u) du,$$

where $C_{k,n}$ and $D_{k,n}$ are universal constants given by Eqs.(8,9).

The reader can observe the analogy with the Laplacian, see e.g. [1]. In particular, one has to distinguish the case of an integrable (resp. non-integrable) singularity for $p_k(\xi)^{-1}$ for the n -dimensional Lebesgue measure.

Proof of Theorem 1. Since $p_k \geq 0$, we define a family $\mathfrak{p}(z)$ of distributions :

$$\forall f \in \mathcal{S}(\mathbb{R}^n) : \langle \mathfrak{p}(z), f \rangle = \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} p_k(\xi)^z \hat{f}(\xi) d\xi.$$

The r.h.s. is holomorphic for $\Re(z) > -1/k$ and, by continuity, we obtain :

$$\lim_{z \rightarrow 0} \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} p_k(\xi)^z \hat{f}(\xi) d\xi = f(0) = \langle \delta, f \rangle. \quad (3)$$

The Laurent development of $\mathfrak{p}$ in $z = -1$ can be written :

$$\mathfrak{p}(z - 1) = \sum_{j=-1}^{-d} z^j \mu_j + \mu_0 + \sum_{j=1}^{\infty} \mu_j z^j. \quad (4)$$

This point is justified in Lemma 2 below. But, according to Eq.(3), we have :

$$\lim_{z \rightarrow 0} \langle \mathfrak{p}(z - 1), P(D)f \rangle = \langle \delta, f \rangle.$$

It is then easy to check that μ_0 is a fundamental solution for P . Also, note that Eqs.(3,4) provide the set of non-trivial relations :

$$P(D)\mu_j = 0, \quad \forall j < 0, \quad (5)$$

in the sense of distributions of $\mathcal{S}'(\mathbb{R}^n)$. The existence of such non-zero μ_j , for $j < 0$, implies the non-uniqueness of fundamental solutions in $\mathcal{S}'(\mathbb{R}^n)$.

Lemma 2 *The distributions $\mathfrak{p}(z - 1)$ are meromorphic on $\mathbb{C}$ with poles located at rational points $z_{j,k} = -\frac{j}{k}$, $j \in \mathbb{N}$.*

Proof. Let $g = \hat{f}$. Using standard polar coordinates we obtain :

$$\langle \mathfrak{p}(z - 1), f \rangle = \frac{1}{(2\pi)^n} \int_{\mathbb{R}_+ \times \mathbb{S}^{n-1}} (r^k p_k(\theta))^{z-1} g(r\theta) r^{n-1} dr d\theta.$$

Here $p_k(\theta)$ is the restriction of p_k to $\mathbb{S}^{n-1}$. Next, if we define :

$$y = (y_1, \dots, y_n) = (r p_k(\theta)^{\frac{1}{k}}, \theta),$$

we obtain a very elementary formulation :

$$\langle \mathfrak{p}(z - 1), f \rangle = \frac{1}{(2\pi)^n} \int_{\mathbb{R}_+} y_1^{k(z-1)} G(y_1) dy_1.$$

This new amplitude G is obtained by pullback and integration :

$$G(y_1) = \int y^*(g(r, \theta) r^{n-1} |Jy|) dy_2 \dots dy_n. \quad (6)$$

We have $G \in \mathcal{S}(\mathbb{R}^+)$ and $G(y_1) = \mathcal{O}(y_1^{n-1})$ in $y_1 = 0$. Starting from :

$$\frac{\partial^k}{\partial y_1^k} y_1^{kz} = \prod_{j=0}^{k-1} (kz - j) y_1^{k(z-1)},$$

after integrations by parts, we accordingly obtain that :

$$\langle \mathfrak{p}(z - 1), f \rangle = \frac{1}{(2\pi)^n} \prod_{j=0}^{k-1} \frac{1}{(kz - j)} \int_{\mathbb{R}_+} y_1^{kz} \partial_{y_1}^k G(y_1) dy_1. \quad (7)$$

The integral of the r.h.s. is holomorphic in the strip $\Re(z) > -\frac{1}{k}$. Finally, we can iterate the previous construction to any order to obtain the result. ■

Since $z = 0$ is a simple pole, the constant term of the Laurent series is :

$$\frac{1}{(2\pi)^n k} \lim_{z \rightarrow 0} \partial_z \left(\prod_{j=1}^{k-1} \frac{1}{(kz - j)} \int_{\mathbb{R}_+} y_1^{kz} \partial_{y_1}^k G(y_1) dy_1 \right).$$

For the calculations, we distinguish integrable and non-integrable singularities.

Case of $k < n$. The derivative of the rational function provides :

$$C_{k,n} = \frac{1}{(2\pi)^n k} (\partial_z (\prod_{j=1}^{k-1} \frac{1}{(kz - j)}))_{z=0} = \frac{(-1)^{k+1}}{(2\pi)^n} \left(\frac{\gamma + \psi(k)}{\Gamma(k)} \right), \quad (8)$$

where γ is Euler's constant and ψ the poly-gamma function of order 0, i.e. :

$$\gamma = \lim_{m \rightarrow \infty} \sum_{j=1}^m \frac{1}{j} - \log(m),$$

$$\psi(z) = \partial_z (\log(\Gamma(z))) = \frac{\Gamma'(z)}{\Gamma(z)}.$$

Accordingly, since G vanishes up to the order $n - 1$ at the origin, we have :

$$C_{k,n} \int_{\mathbb{R}_+} \partial_{y_1}^k G(y_1) dy_1 = -C_{k,n} \partial_{y_1}^{k-1} G(0) = 0.$$

On the other side, by derivation of the integral, we find the term :

$$\frac{(-1)^{k-1}}{(k-1)!} \int_{\mathbb{R}_+} \log(y_1) \partial_{y_1}^k G(y_1) dy_1.$$

Since $k < n$, we can integrate by parts to finally obtain :

$$\int_{\mathbb{R}_+} G(y_1) \frac{dy_1}{y_1^k} = \int_{\mathbb{R}_+ \times \mathbb{S}^{n-1}} p_k(\theta)^{-1} g(r\theta) r^{n-1-k} dr d\theta.$$

Here, we have replaced the expression for our amplitude, via inversion of our diffeomorphism. This proves the first statement of Theorem 1. ■

Case of $k \geq n$. The term attached to the derivative of the rational function is non-zero and provides :

$$-C_{k,n} \partial_{y_1}^{k-1} G(0) + \frac{1}{(2\pi)^n} \frac{(-1)^{k-1}}{(k-1)!} \int_{\mathbb{R}_+} \log(y_1) \partial_{y_1}^k G(y_1) dy_1.$$

We compute first the derivative of G . Contrary to the case $k < n$ we cannot take the limit directly but we will reach the result with the Schwartz kernel technic. By Fourier inversion formula we have :

$$\partial_{y_1}^{k-1} G(0) = \frac{1}{2\pi} \int e^{-i\xi u} (i\xi)^{k-1} G(u) du d\xi.$$

We can extend the integral w.r.t. u on the whole line by extending G by zero for $u < 0$. Going back to initial coordinates provides :

$$\partial_{y_1}^{k-1} G(0) = \frac{1}{2\pi} \int e^{-i\xi r} (i\xi)^{k-1} r^{n-1} \int_{\mathbb{S}^{n-1}} g(r\theta) p_k(\theta)^{-1} d\theta dr d\xi.$$

The r.h.s. is exactly the derivative of order $k-1$ of $A(g)$ defined in Eq.(2). By the same technic, the logarithmic integral gives the non-local contribution :

$$\frac{1}{(k-1)!} \int_{\mathbb{R}_+} \log(u) \frac{\partial^k A(g)}{\partial r^k}(u) du.$$

In particular the second universal constant of Theorem 1 is :

$$D_{k,n} = \frac{1}{(2\pi)^n} \frac{(-1)^{k-1}}{(k-1)!}. \quad (9)$$

This proves the second assertion of Theorem 1. ■

The main result holds also holds for elliptic pseudo-differential operators with homogeneous symbol of degree $\alpha \in]0, \infty[$. Using $k = E(\alpha) + 1$ integrations by parts in Eq.(7), where $E(x)$ is the integer part of x , the construction remains the same and all formulas can be analytically continued w.r.t. the degree $\alpha > 0$. This continuation is trivial since the functions of Eqs.(8,9), defining respectively $C_{n,k}$ and $D_{n,k}$, are analytic w.r.t. the variable $k > 0$.

References

- [1] L.Hörmander, The analysis of linear partial differential operators vol.1, Distribution theory and Fourier analysis. Springer-Verlag, Berlin (1983).
- [2] L.Hörmander, The analysis of linear partial differential operators vol.2, Differential operators with constant coefficients. Springer-Verlag, Berlin (1983).
- [3] P.Wagner, Fundamental solutions of real homogeneous cubic operators of principal type in three dimensions, Acta Mathematica **182** (1999) no.2, 283-300.
- [4] P.Wagner, On the fundamental solutions of a class of elliptic quartic operators in dimension 3, Journal de Mathématiques Pures et Appliquées, Neuvième Série **81** (2002) no. 11, 1191-1206.